

Brown Dwarfs, White Knights and Demons

Geza Gyuk

S.I.S.S.A., via Beirut 2-4, 34014 Trieste, Italy

and

N. Wyn Evans

Theoretical Physics, Department of Physics, 1 Keble Rd, Oxford, OX1 3NP, UK

and

Evalyn I. Gates

Departments of Astronomy & Astrophysics, The University of Chicago, Chicago, IL 60637

Adler Planetarium, 1300 Lake Shore Drive, Chicago, IL 60605

Received _____; accepted _____

ABSTRACT

This paper investigates the hypothesis that the lensing objects towards the Large Magellanic Cloud (LMC) are brown dwarfs by analysing the effects of velocity anisotropy on the inferred microlensing masses. To reduce the masses, the transverse velocity of the lenses with respect to the microlensing tube must be minimised. In the outer halo, radial anisotropy is best for doing this; closer to the solar circle, azimuthal anisotropy is best. By using a constraint on the total kinetic energy of the tracer population from the Jeans equations, the microlensing mass is minimised over orientations of the velocity dispersion tensor. This minimum mass is $\approx 0.1 M_{\odot}$, which lies above the hydrogen burning limit. This demonstrates explicitly that populations of brown dwarfs with smoothly decreasing densities and dynamically mixed velocity distributions cannot be responsible for the microlensing events. Brown dwarfs are no white knights! There is one caveat. If there are demons sitting on the microlensing tube, they can drop brown dwarfs so as to reproduce the microlensing data-set exactly. Such a distribution is not smooth and does not give well-mixed velocities in phase space. It is a permissible solution only if the outer halo is dynamically young and lumpy. In such a case, theorists cannot rule out brown dwarfs. Only exorcists can!

Subject headings: Galaxy: halo – Galaxy: kinematics and dynamics – gravitational lensing – dark matter

1. INTRODUCTION

The MACHO collaboration has interpreted its observations of microlensing events towards the Large Magellanic Cloud (LMC) as evidence that about one third of the halo of our own halo exists in the form of objects of around 0.5 solar mass (Alcock et al. 1997). Unfortunately, there are seemingly insuperable objections to all the obvious candidates for the lensing population. Normal stars would be visible (Alcock et al. 1997), white dwarfs are ruled out by current population II abundance ratios (Fields, Mathews & Schramm 1997, Gibson & Mould 1997), while the Hubble Deep Field gives stringent restrictions on the contribution of red dwarfs (Graff & Freese 1996). The microlensing events would be easier to understand if the characteristic mass of the lensing objects was below the hydrogen-burning limit ($\approx 0.08M_{\odot}$). Of course, a lensing population of brown dwarfs would be much too dark to be visible and there is no conflict either with the metallicity data or the Hubble Deep Field star counts. So, it is natural to ask the questions: Can the deflectors be brown dwarfs? Is it possible that the masses of the microlenses have hitherto been overestimated? The aim of this Letter is to answer these questions.

Uncertainties in estimates of the lens candidates arise from two fundamental sources: low number statistics and modelling error. Although the number of microlensing events observed towards the LMC is still low, a determination of the average mass for a *given* halo model can be obtained with perhaps 50% accuracy (see eg. Mao & Paczyński 1996; Alcock et al. 1997). This number is expected to improve substantially over the course of the next few years as new events are detected. A much more important source of error comes from our ignorance of the structure of the outer Milky Way halo. The halo models used by Alcock et al. (1997) are either isotropic, such as the cored isothermal sphere (Griest 1991), or they are very nearly so, such as the power-law models (Evans 1994). Alcock et al. (1997; see especially Figures 17 and 24) plot likelihood contours in the plane of the lens mass

and baryon fraction of the halo. The striking elongation of the contours along the baryon fraction axis suggests that there is comparatively little uncertainty in the mass estimates of the microlenses for a given model. One worry is that this propitious state of affairs is a consequence of using halo models that all pretty much look the same! In a percipient investigation, Marković & Sommer-Larsen (1997) looked at a wider range of halos, including some with anisotropic velocity distributions. They found that ~ 100 events (an order of magnitude more than presently available) are needed to estimate the average mass. This large error bar includes both the modelling and the statistical uncertainty. The claims of Marković & Sommer-Larsen (1997) may be somewhat overstated because of the uniform priors used in their Monte Carlo simulations. However, Mao & Paczyński (1996) have also emphasised the difficulty of drawing firm conclusions about the mass distribution of the lenses from the limited sample available. These two papers consider both the statistical and the modelling uncertainties together. The focus of our paper is on the modelling uncertainty alone. Our aim is to demonstrate unambiguously that the modelling uncertainties cannot be responsible for the high average mass estimates of Alcock et al. (1997).

Low mass lenses such as brown dwarfs are already ruled only for halo models with negligible rotation and isotropic velocity dispersions (e.g., Chabrier, Segretain & Méra 1996). To rule out the hypothesis that the lenses are brown dwarfs requires a thorough investigation of halo models with very different kinematics – in particular with different streaming velocities and different random motions. Gyuk & Gates (1998) have already shown that rotating halos are unable to reduce the microlensing mass estimates below about $0.25 M_{\odot}$ (unless all the lensing takes place very close to the Sun). This Letter will examine the effects of anisotropy and show that the associated *modeling uncertainties* cannot cause the high lens mass estimates.

2. VELOCITY ANISOTROPY AND THE MINIMUM MASS OF THE MICROLENSING OBJECTS

Let us start with a thought experiment. Suppose a stationary observer views a stationary source through a population of lenses with density ρ . The timescale of any lensing event is related to the Einstein radius R_E and the transverse velocity v_T by

$$t_0 = \frac{R_E}{v_T} = \frac{1}{v_T} \sqrt{\frac{4GM D_d (D_s - D_d)}{c^2 D_s}}, \quad (1)$$

where M is the mass of the lens and D_d and D_s are the distances to deflector and source. Suppose now that the distribution of transverse velocities of the lenses is Gaussian with a dispersion σ_T . The microlensing optical depth τ is well-known to be independent of the masses of the lenses (Press & Gunn 1973). The rate of microlensing Γ is (e.g., Griest 1991)

$$\Gamma = (2\pi)^{1/2} \frac{\sigma_T}{M^{1/2}} \int_0^{D_s} dD_d \rho(D_d) \sqrt{\frac{4GD_d(D_s - D_d)}{c^2 D_s}}, \quad (2)$$

and the timescale histogram is

$$\frac{d\Gamma}{dt_0} = \frac{8A^2 \sigma_T^2}{M} \int_0^{D_s} dD_d \rho(D_d) D_d^2 (D_s - D_d)^2 \exp[-AD_d(D_s - D_d)], \quad (3)$$

with $A = 2GM/(D_s c^2 t_0 \sigma_T^2)$. This demonstrates explicitly that all the microlensing quantities ($\tau, \Gamma, \frac{d\Gamma}{dt_0}$) depend only on the ratio M/σ_T^2 . Given the microlensing data-set alone, it is not possible to constrain the mass of the deflectors at all! Any mass estimate is solely a consequence of assumptions regarding the transverse velocity dispersions. The same data will be consistent with smaller inferred mass if the transverse motions are reduced. This degeneracy between mass and velocity can be partially lifted if parallax effects (Refsdal 1966, Griest 1991, Gould 1994), or finite source size effects (Nemiroff 1997) can be detected in the lightcurve.

Of course, the analysis of the microlensing events towards the LMC is more complex than this thought experiment. Both the Sun and the LMC are moving and therefore the

expectation value of the transverse velocity of the lens with respect to the microlensing tube cannot be made arbitrarily small just by changing the velocity anisotropy of the lenses. Figure 1 shows a planform of the Sun and the LMC projected on the Galactic equatorial plane. The line of sight from the Sun to the LMC is shown as a dashed line. In the outer parts of the halo, this line of sight is aligned very nearly with the radial direction of the spherical polar coordinate systems. Radial anisotropy of the velocity dispersion tensor is the best option for reducing the mass estimates. Nearer the solar circle, the offset of the Sun from the Galactic Center becomes important. The line of sight is aligned more nearly with the azimuthal direction. This means that radial anisotropy is now dangerous. The best recipe for the minimum microlensing mass is to allow the velocity dispersion tensor to be azimuthally distended near the Sun and to become radially distended in the outer halo.

As a simple model, let us assume that the density of the lensing population is smooth and falls off like a power of the distance ($\rho \propto r^{-\gamma}$). We take the overall potential to be a power-law model, so that the circular velocity, v_{circ} , falls like $r^{-\beta/2}$. Rich families of solutions to the Jeans equations for power-law density distributions in power-law potentials are known (Evans, Häfner & de Zeeuw 1997). These are all aligned in the spherical polar coordinates, but vary in the anisotropy of the principal components of the velocity dispersion tensor σ_i . The detailed Jeans solutions all satisfy the constraint (see eq. (3.8) of Evans, Häfner & de Zeeuw 1997)

$$\sum_{i=1}^3 \sigma_i^2 \gtrsim v_{\text{circ}}^2 \times \min\left(1, \frac{1}{\beta + \gamma - 2}\right). \quad (4)$$

From the standpoint of minimising the microlensing mass estimates, the best of all possible worlds is to replace the inequality in the above expression with an equality. This means that the total kinetic energy required to support the lensing population against gravity is underestimated. The inferred microlensing mass will always be lower than the true mass. We allow the ratio of the principal components of the velocity dispersions to vary subject

only to the condition that the sum of the components does not violate the inequality (4). Thus, the Jeans equations are not satisfied spot-wise, but only in a gross sense. The total kinetic energy cannot be reduced further without violating the rules of gravitational physics. If all the deflectors are $1M_{\odot}$ objects, the rate is (see e.g., Griest 1991, Gyuk & Gates 1998)

$$\Gamma = 2 \int_0^{D_s} dD_d R_E(D_d) \rho(D_d) \langle |v_{\perp}| \rangle. \quad (5)$$

Here $\langle |v_{\perp}| \rangle$ is the average value of the transverse velocity of the lens with respect to the microlensing tube. The best estimator of the average event duration $\langle t_e \rangle$ is 61 days (see Appendix A of Gyuk & Gates 1998). This uses the events and the efficiencies given in Alcock et al. (1997a). So the microlensing mass estimate $m_{\min}$ is

$$m_{\min} \gtrsim \left[\langle t_e \rangle \frac{\Gamma}{\tau} \right]^2. \quad (6)$$

Our algorithm for finding the minimum microlensing mass estimate is as follows. Choose the model parameters β and γ and an alignment of the velocity dispersion tensor, and then apportion the total kinetic energy into the three principal components subject only to the constraint (4). The velocity anisotropy may be parametrised by

$$\lambda = \frac{\sigma_2^2 + \sigma_3^2}{2\sigma_1^2}, \quad \mu = \frac{\sigma_3^2}{\sigma_2^2}. \quad (7)$$

Let us insist that the velocity ellipsoid cannot be anisotropic by more than a 4 : 1 ratio; that is to say, λ and μ must lie within the range 1/16 to 16. For comparison, Freeman (1987) reports that the Population II stars in the spheroid have velocity dispersions at the solar position oriented on the cylindrical polar coordinate system such that $(\sigma_R, \sigma_{\phi}, \sigma_z) \approx (140, 100, 75) \text{ kms}^{-1}$. The microlensing mass is to be minimised as both the anisotropy parameters λ and μ and the alignment of the velocity ellipsoid are varied.

After a little thought, it is obvious what the alignment is for the minimum mass estimate – the best of all possible worlds is when the velocity ellipsoid is aligned along the

microlensing tube itself, with as much kinetic energy as possible put into motions along the tube and as little as possible put into transverse motions. Figure 2 shows the inferred mass as a function of γ where the mass has been minimized by allowing λ and μ to float over their respective ranges. Different curves correspond to the range $-0.25 \leq \beta \leq 0.25$. In such a situation, λ always prefers to be as low as possible, while μ is only weakly constrained. The minimum mass estimate ranges from $0.1 M_{\odot}$ when $\gamma = 2.0$ to $0.25 M_{\odot}$ when $\gamma = 4.0$. The inferred mass is always larger than the hydrogen-burning limit. This leads us to the main result of the Letter. *If the density of the microlensing population is smooth and monotonic decreasing (that is, reasonably well-approximated by a power-law), then the microlenses cannot be brown dwarfs, irrespective of the details of their kinematics.* The strength of this statement is that it is based on the Jeans equations and therefore robust.

3. A MODEST ESTIMATE

Let us emphasise that this algorithm for obtaining the minimum mass gives a value that is *very much a lower limit*. It uses a number of gratuitous approximations, all of which act to reduce the mass estimate. For example, the Jeans solutions of reasonable tracer populations may possess a kinetic energy greater than the minimum prescribed by eq. (4). Again, almost certainly, the alignment along the microlensing tube that yields the minimum mass cannot be built – that is, there is no set of stellar orbits that can be superposed to yield a true dynamical model corresponding to the Jeans solution. Making the model more realistic will necessarily require more massive lenses. In this section we provide an estimate of the more modest reduction in the microlensing masses expected from velocity anisotropy for one particular reasonably realistic model of the halo.

To do this, let us build Jeans solutions of tracer populations with the density of the

Jaffe (1983) sphere

$$\rho = \frac{M}{4\pi r_J^3} \frac{r_J^4}{r^2(r + r_J)^2} \quad (8)$$

in a spherical isothermal halo potential. Here, r_J is a scale-length that describes when the density turns over. A typical estimate of its value might be $r_J \sim 50\text{kpc}$ (Kochanek 1996, Wilkinson & Evans 1998). Let the anisotropy be defined as

$$\frac{\sigma_\theta^2}{\sigma_r^2} = \frac{\sigma_\phi^2}{\sigma_r^2} = \lambda_\infty + \frac{r_\lambda}{r}. \quad (9)$$

This simple ansatz allows the kinematics to change from azimuthal anisotropy to radial anisotropy or vice versa. Here, λ_∞ is the value of the anisotropy at infinity, whereas r_λ is a scale-length on which the anisotropy changes. The solution of the spherical Jeans equation is readily found by the method of integrating factors as (see Binney & Tremaine 1987)

$$\sigma_r^2 = v_{\text{circ}}^2 \exp(-2r_\lambda/r) r^{2\lambda_\infty} (r + r_J)^2 \int_r^\infty \frac{dr \exp(2r_\lambda/r)}{r^{2\lambda_\infty+1} (r + r_J)^2}. \quad (10)$$

The azimuthal dispersions now follow from (9). The isotropic model has $\lambda_\infty = 1$ and $r_\lambda = 0$ and a mass estimate (given by eq. 6) of $0.348 M_\odot$. This can be reduced by anisotropy. The microlensing mass estimate in the plane of the asymptotic anisotropy λ_∞ and the anisotropy scale r_λ is shown in Figure 3. For this set of Jeans solutions – in which the anisotropy can change significantly but not dramatically – there is no hope of using anisotropy by itself to reduce the microlensing mass estimate below $\approx 0.3 M_\odot$.

4. DISCUSSION AND CONCLUSIONS

If the density of the microlenses is smooth and decreasing, then they cannot be brown dwarfs. This holds irrespective of the details of their kinematics. This general result follows because the Jeans equations (or, equivalently, the virial theorem) imply the existence of an irreducible minimum kinetic energy to support the lensing population against gravity. Even

in the optimum alignment of the velocity dispersion tensor of the lenses, this must yield sufficient transverse motion so that the minimum mass is $\approx 0.1 M_{\odot}$ for halo models with flat rotation curves. This is above the hydrogen burning limit.

There is a way to save brown dwarfs. Let us imagine a collection of demons sitting on the microlensing tube. One of the demons at a heliocentric distance of 20 kpc launches a brown dwarf of mass $0.06 M_{\odot}$ with a velocity of 106 km s^{-1} across our line of sight ... and this causes event # 4 with a blended timescale of 39.5 days. A second demon sitting on the microlensing tube at 30 kpc lobs a brown dwarf with a velocity of just 75 km s^{-1} ... and this gives event # 5 with a blended timescale of 55.5 days, and so on. Of course, demons can exactly reproduce the dataset reported by Alcock et al. (1997) by dropping brown dwarfs from the microlensing tube. The density of brown dwarfs so produced is neither spherical nor axisymmetric nor in a steady-state. The velocity distribution is not dynamically well-mixed and the time averages theorem (Binney & Tremaine 1987, p. 171), which is the fundamental result underpinning steady-state stellar dynamics, does not hold. If it did, we could infer the existence of further brown dwarfs at different phases of the same orbits and show that they produce microlensing events that are not seen. Such a model is possible if the halo is very blobby (e.g. Lynden-Bell 1994, Lynden-Bell & Lynden-Bell 1995). Then, in every direction that one looks (including $\ell = 280^{\circ}, b = -33^{\circ}$), there may be garbage heaps of brown dwarfs whose density and velocity distributions are lumpy. This possibility cannot be ruled out from the microlensing data-set alone.

Most of this work was done during a visit to SISSA, Trieste. NWE wishes to thank the Astrophysics Sector in general, and Dennis Sciama and John Miller in particular, for their kindness and their hospitality. NWE thanks James Binney for numerous insightful remarks on this subject.

REFERENCES

- Alard C., 1997. AA, 321, 424
- Alcock C., et al. 1997, ApJ, 486, 697
- Binney J., Tremaine S. 1987, Galactic Dynamics (Princeton University Press, Princeton)
- Chabrier G., Ségretain L., Méra D., 1996, ApJ, 468, L21
- Evans N.W. 1994, MNRAS, 267, 333
- Evans N.W., Häfner R., de Zeeuw P.T. 1997, MNRAS, 286, 315
- Fields B., Mathews G.J., Schramm D.N. 1997, ApJ, 483, 625
- Freeman K. 1987, ARAA, 25, 603
- Gates E., Gyuk G., Turner M.S. 1995, ApJ, L123
- Gibson B., Mould J. 1997, ApJ, 482, 98.
- Gould A. 1994, ApJ, 421, L75
- Graff D., Freese K. 1996, ApJ, 456, L49
- Griest K. 1991, ApJ, 366, 412
- Gyuk G., Gates E. 1998, MNRAS, 294, 682
- Jaffe W. 1983, MNRAS, 202, 995
- Kochanek C. 1996, ApJ, 457, 228
- Lynden-Bell D. 1994, In: Dwarf Galaxies, eds G. Meylan, P. Prugniel, ESO Publications,
p. 548

Lynden-Bell D., Lynden-Bell R. 1995, MNRAS, 275, 439

Mao S., Paczyński B. 1996, ApJ, 473, 57

Marković D., Sommer-Larsen J. 1997, MNRAS, 288, 733

Nemiroff R. 1997, ApJ, 486, 693

Press W., Gunn J. 1973, ApJ, 185, 397

Refsdal S. 1966, MNRAS, 134, 315

di Stefano R., Esin A.A. 1995, ApJ, 448, L1

Wilkinson M., Evans N.W. 1998, in preparation




Fig. 1.— The positions of the Sun and the Large Magellanic Cloud (LMC) projected onto the Galactic Plane. The line of sight from the Sun to the LMC is marked. This is almost radially aligned in the outer halo, but azimuthally aligned near the Solar circle.




Fig. 2.— The minimum mass estimate as a function of the steepness of the lens density distribution, γ . As described in the text both anisotropies have been allowed to float. The break in the curve at $\gamma = 3$ is due to the prescription for finding the minimum kinetic energy. For all γ the microlensing mass is $\gtrsim 0.1 M_{\odot}$, which is too massive for cold, degenerate brown dwarfs. While recognising the statistical uncertainties are still great, the virtue of this figure is that it demonstrates that the modelling uncertainties associated with anisotropy cannot be responsible for the high lens mass estimates.




Fig. 3.— Contour plots of the minimum microlensing mass for the Jaffe model in the plane of the model parameters $(r_\lambda, \log \lambda_\infty)$. The contours are spaced at intervals of $0.01 M_\odot$. Anisotropy does change the mass estimate – but never below $\approx 0.3 M_\odot$. For this particular Jaffe model with $r_J \sim 50$ kpc, radial anisotropy is best for reducing the mass estimate.